

Common Medications that if Combined have Potential for Negative Interactions

DISCLAIMER: *The information contained herein should NOT be used as a substitute for the advice of an appropriately qualified and licensed physician or other health care provider. list of medications is not all-inclusive, i.e.: there are many other common and uncommon medications in existence. Additionally, the interactions listed next to each medication do not entail all possible interactions that may occur.*

USE “CTRL” + “SHIFT” + “F” to Search for Drug(s)

DO NOT EVER USE TOGETHER

Never use this combination of drugs because of high risk for dangerous interaction.

Ziprasidone + Escitalopram

Ziprasidone and Escitalopram both increase causing a dangerous abnormal heart rhythm.

Ziprasidone + Citalopram

Ziprasidone and Citalopram both increase causing a dangerous abnormal heart rhythm.

Additional Information: Caution; use of these drugs may cause heart beat changes; ECG (cardiac) monitoring by your doctor is recommended.

Epinephrine + Lurasidone

Epinephrine increases toxicity of Lurasidone by added drug effects.

Additional Information: Interaction applies only when there is an acute overdose of lurasidone. Epinephrine may enhance hypotensive (abnormally low blood pressure) effects of lurasidone when overdosed.

POTENTIAL FOR SERIOUS NEGATIVE INTERACTIONS

These combinations of drugs have high risk for serious and dangerous interaction. Potential for serious interaction; regular monitoring by your doctor required or alternate medication may be needed.

Hydroxyzine + Trazodone

Hydroxyzine and Trazodone both increase causing a dangerous abnormal heart rhythm.

Hydroxyzine + Paliperidone

Hydroxyzine and Paliperidone both increase causing a dangerous abnormal heart rhythm.

Hydroxyzine + Quetiapine

Hydroxyzine increases toxicity of Quetiapine by causing a dangerous abnormal heart rhythm.

Additional Information: This combination increases the risk of abnormal heartbeats.

Hydroxyzine + Ziprasidone

Hydroxyzine increases toxicity of Ziprasidone by causing a dangerous abnormal heart rhythm.

Additional Information: This combination increases the risk of abnormal heartbeats.

Hydroxyzine + Fluoxetine

Hydroxyzine increases toxicity of Fluoxetine by causing a dangerous abnormal heart rhythm.

Additional Information: This combination increases the risk of abnormal heartbeats.

Hydroxyzine + Citalopram

Hydroxyzine increases toxicity of Citalopram by causing a dangerous abnormal heart rhythm.

Additional Information: This combination increases the risk of abnormal heartbeats.

Hydroxyzine + Iloperidone

Hydroxyzine increases toxicity of Iloperidone by causing a dangerous abnormal heart rhythm.

Additional Information: This combination increases the risk of abnormal heartbeats.

Clonidine + Tramadol

Clonidine , Tramadol . Either increases toxicity of the other by added drug effects.

Topiramate + Ethinylestradiol

Topiramate will decrease the level or effect of Ethinylestradiol by altering drug metabolism.

Additional Information: The effect of hormonal contraceptives may be reduced. Use of a nonhormonal contraceptive is recommended.

Topiramate + Norethindrone Acetate

Topiramate will decrease the level or effect of Norethindrone Acetate by altering drug metabolism.

Additional Information: This combination reduces the effect of the hormonal contraceptive and may increase incidence of menstruation associated adverse effects.

Topiramate + Norethindrone

Topiramate will decrease the level or effect of Norethindrone by altering drug metabolism.

Additional Information: This combination reduces the effect of the hormonal contraceptive and may increase incidence of menstruation associated adverse effects.

Citalopram + Ziprasidone

Citalopram and Ziprasidone both increase causing a dangerous abnormal heart rhythm.

Citalopram + Amitriptyline

Citalopram and Amitriptyline both increase causing a dangerous abnormal heart rhythm.

Citalopram and Amitriptyline both increase affecting serotonin levels in the blood. Too much serotonin is a potentially life-threatening situation. Severe signs and symptoms include high blood pressure and increased heart rate that lead to shock.

Additional Information: Risk of serious adverse effects may increase including serotonin syndrome or neuroleptic malignant syndrome (a neurological disorder). Potential risk for QT prolongation (abnormal heart beats). ECG (cardiac) monitoring by your doctor is recommended.

Citalopram and Amitriptyline both increase causing a dangerous abnormal heart rhythm.

Citalopram and Amitriptyline both increase affecting serotonin levels in the blood. Too much serotonin is a potentially life-threatening situation. Severe signs and symptoms include high blood pressure and increased heart rate that lead to shock.

Additional Information: Risk of serious adverse effects may increase including serotonin syndrome or neuroleptic malignant syndrome (a neurological disorder). Potential risk for QT prolongation (abnormal heart beats). ECG (cardiac) monitoring by your doctor is recommended.

Citalopram + Trazodone

Citalopram and Trazodone both increase causing a dangerous abnormal heart rhythm.

Citalopram + Risperidone

Citalopram and Risperidone both increase causing a dangerous abnormal heart rhythm.

Citalopram + Fluoxetine

Citalopram and Fluoxetine both increase affecting serotonin levels in the blood. Too much serotonin is a potentially life-threatening situation. Severe signs and symptoms include high blood pressure and increased heart rate that lead to shock.

Additional Information: Combination may increase risk of potentially life-threatening reactions such as serotonin syndrome or neuroleptic malignant syndrome (a neurological disorder).

Citalopram + Venlafaxine

Citalopram and Venlafaxine both increase affecting serotonin levels in the blood. Too much serotonin is a potentially life-threatening situation. Severe signs and symptoms include high blood pressure and increased heart rate that lead to shock.

Additional Information: Combination may increase risk of potentially life-threatening reactions such as serotonin syndrome or neuroleptic malignant syndrome (a neurological disorder).

Citalopram + Escitalopram

Citalopram and Escitalopram both increase affecting serotonin levels in the blood. Too much serotonin is a potentially life-threatening situation. Severe signs and symptoms include high blood pressure and increased heart rate that lead to shock.

Additional Information: Combination may increase risk of potentially life-threatening reactions such as serotonin syndrome or neuroleptic malignant syndrome (a neurological disorder).

Citalopram + Buspirone

Citalopram and Buspirone both increase affecting serotonin levels in the blood. Too much serotonin is a potentially life-threatening situation. Severe signs and symptoms include high blood pressure and increased heart rate that lead to shock.

Additional Information: Combination may increase risk of potentially life-threatening reactions such as serotonin syndrome or neuroleptic malignant syndrome (a neurological disorder).

Fluoxetine + Bupropion

Fluoxetine increases toxicity of Bupropion by unspecified interaction mechanism.

Additional Information: Keep bupropion dose as low as possible as seizures may occur.

Fluoxetine + Buspirone

Fluoxetine and Buspirone both increase affecting serotonin levels in the blood. Too much serotonin is a potentially life-threatening situation. Severe signs and symptoms include high blood pressure and increased heart rate that led to shock.

Fluoxetine + Trazodone

Fluoxetine and Trazodone both increase affecting serotonin levels in the blood. Too much serotonin is a potentially life-threatening situation. Severe signs and symptoms include high blood pressure and increased heart rate that lead to shock.

Fluoxetine + Amitriptyline

Fluoxetine and Amitriptyline both increase affecting serotonin levels in the blood. Too much serotonin is a potentially life-threatening situation. Severe signs and symptoms include high blood pressure and increased heart rate that lead to shock.

Fluoxetine will increase the level or effect of Amitriptyline by altering drug metabolism.

Fluoxetine and Amitriptyline both increase affecting serotonin levels in the blood. Too much serotonin is a potentially life-threatening situation. Severe signs and symptoms include high blood pressure and increased heart rate that lead to shock.

Fluoxetine will increase the level or effect of Amitriptyline by altering drug metabolism.

Fluoxetine + Venlafaxine

Fluoxetine and Venlafaxine both increase affecting serotonin levels in the blood. Too much serotonin is a potentially life-threatening situation. Severe signs and symptoms include high blood pressure and increased heart rate that lead to shock.

Fluoxetine + Iloperidone

Fluoxetine will increase the level or effect of Iloperidone by altering drug metabolism.

Fluoxetine + Aripiprazole

Fluoxetine will increase the level or effect of Aripiprazole by altering drug metabolism.

Fluoxetine and Aripiprazole both increase causing a dangerous abnormal heart rhythm.

Fluoxetine + Risperidone

Fluoxetine will increase the level or effect of Risperidone by altering drug metabolism.

Fluoxetine + Escitalopram

Fluoxetine will increase the level or effect of Escitalopram by altering drug metabolism.

Fluoxetine and Escitalopram both increase affecting serotonin levels in the blood. Too much serotonin is a potentially life-threatening situation. Severe signs and symptoms include high blood pressure and increased heart rate that lead to shock.

Fluvoxamine + Melatonin

Fluvoxamine will increase the level or effect of Melatonin by altering drug metabolism.

Fluvoxamine + Omeprazole

Fluvoxamine will increase the level or effect of Omeprazole by altering drug metabolism.

Fluvoxamine + Trazodone

Fluvoxamine and Trazodone both increase affecting serotonin levels in the blood. Too much serotonin is a potentially life-threatening situation. Severe signs and symptoms include high blood pressure and increased heart rate that lead to shock.

Fluvoxamine + Buspirone

Fluvoxamine and Buspirone both increase affecting serotonin levels in the blood. Too much serotonin is a potentially life-threatening situation. Severe signs and symptoms include high blood pressure and increased heart rate that lead to shock.

Fluvoxamine + Escitalopram

Fluvoxamine and Escitalopram both increase affecting serotonin levels in the blood. Too much serotonin is a potentially life-threatening situation. Severe signs and symptoms include high blood pressure and increased heart rate that lead to shock.

Fluvoxamine + Venlafaxine

Fluvoxamine and Venlafaxine both increase affecting serotonin levels in the blood. Too much serotonin is a potentially life-threatening situation. Severe signs and symptoms include high blood pressure and increased heart rate that lead to shock.

Fluvoxamine + Fluoxetine

Fluvoxamine and Fluoxetine both increase affecting serotonin levels in the blood. Too much serotonin is a potentially life-threatening situation. Severe signs and symptoms include high blood pressure and increased heart rate that lead to shock.

Fluvoxamine + Citalopram

Fluvoxamine and Citalopram both increase affecting serotonin levels in the blood. Too much serotonin is a potentially life-threatening situation. Severe signs and symptoms include high blood pressure and increased heart rate that lead to shock.

Fluvoxamine + Amitriptyline

Fluvoxamine and Amitriptyline both increase affecting serotonin levels in the blood. Too much serotonin is a potentially life-threatening situation. Severe signs and symptoms include high blood pressure and increased heart rate that lead to shock.

Fluvoxamine + Bupropion

Fluvoxamine increases toxicity of Bupropion by Other mechanism.

Venlafaxine + Bupropion

Venlafaxine increases toxicity of Bupropion by unspecified interaction mechanism.

Additional Information: Keep bupropion dose as low as possible as seizures may occur.

Venlafaxine + Buspirone

Venlafaxine and Buspirone both increase affecting serotonin levels in the blood. Too much serotonin is a potentially life-threatening situation. Severe signs and symptoms include high blood pressure and increased heart rate that lead to shock.

Venlafaxine + Amitriptyline

Venlafaxine and Amitriptyline both increase affecting serotonin levels in the blood. Too much serotonin is a potentially life-threatening situation. Severe signs and symptoms include high blood pressure and increased heart rate that lead to shock.

Venlafaxine + Pseudoephedrine

Venlafaxine increases effects of Pseudoephedrine by adrenaline effects, which affect blood pressure and heart rate.

Amitriptyline + Ziprasidone

Amitriptyline and Ziprasidone both increase causing a dangerous abnormal heart rhythm.

Amitriptyline + Trazodone

Amitriptyline and Trazodone both increase causing a dangerous abnormal heart rhythm.

Amitriptyline and Trazodone both increase affecting serotonin levels in the blood. Too much serotonin is a potentially life-threatening situation. Severe signs and symptoms include high blood pressure and increased heart rate that lead to shock.

Amitriptyline + Pseudoephedrine

Amitriptyline increases effects of Pseudoephedrine by adrenaline effects, which affect blood pressure and heart rate.

Amitriptyline + Buspirone

Amitriptyline and Buspirone both increase affecting serotonin levels in the blood. Too much serotonin is a potentially life-threatening situation. Severe signs and symptoms include high blood pressure and increased heart rate that lead to shock.

Amitriptyline + Dexmethylphenidate

Amitriptyline , Dexmethylphenidate Other mechanism.

Additional Information: Tricyclic antidepressants may increase or decrease the effect of the other drug when taken in this combination.

Amitriptyline + Dextroamphetamine

Amitriptyline , Dextroamphetamine Other mechanism.

Additional Information: Tricyclic antidepressants may increase or decrease the effect of the other drug when taken in this combination.

Amitriptyline + Albuterol

Amitriptyline , Albuterol Other mechanism.

Additional Information: Tricyclic antidepressants may increase or decrease the effect of the other drug when taken in this combination.

Amitriptyline + Lisdexamfetamine

Amitriptyline , Lisdexamfetamine Other mechanism.

Additional Information: Tricyclic antidepressants may increase or decrease the effect of the other drug when taken in this combination.

Amitriptyline + Guanfacine

Amitriptyline decreases effects of Guanfacine by Other mechanism.

Additional Information: One drug reduces the effect of the other.

Amitriptyline + Epinephrine

Amitriptyline , Epinephrine Other mechanism.

Additional Information: Tricyclic antidepressants may increase or decrease the effect of the other drug when taken in this combination.

Amitriptyline and Epinephrine both increase causing a dangerous abnormal heart rhythm.

Amitriptyline + Clonidine

Amitriptyline decreases effects of Clonidine by Other mechanism.

Additional Information: One drug reduces the effect of the other.

Trazodone + Clonidine

Trazodone decreases effects of Clonidine by Other mechanism.

Additional Information: One drug reduces the effect of the other.

Trazodone + Epinephrine

Trazodone , Epinephrine Other mechanism.

Additional Information: Tricyclic antidepressants may increase or decrease the effect of the other drug when taken in this combination.

Trazodone and Epinephrine both increase causing a dangerous abnormal heart rhythm.

Trazodone + Guanfacine

Trazodone decreases effects of Guanfacine by Other mechanism.

Additional Information: One drug reduces the effect of the other.

Trazodone + Lisdexamfetamine

Trazodone , Lisdexamfetamine Other mechanism.

Additional Information: Tricyclic antidepressants may increase or decrease the effect of the other drug when taken in this combination.

Trazodone + Albuterol

Trazodone , Albuterol Other mechanism.

Additional Information: Tricyclic antidepressants may increase or decrease the effect of the other drug when taken in this combination.

Trazodone + Dextroamphetamine

Trazodone , Dextroamphetamine Other mechanism.

Additional Information: Tricyclic antidepressants may increase or decrease the effect of the other drug when taken in this combination.

Trazodone + Bupropion

Trazodone increases toxicity of Bupropion by unspecified interaction mechanism.

Additional Information: Keep Bupropion dose as low as possible as seizure may occur.

Trazodone + Dexmethylphenidate

Trazodone , Dexmethylphenidate Other mechanism.

Additional Information: Tricyclic antidepressants may increase or decrease the effect of the other drug when take in this combination.

Trazodone + Pseudoephedrine

Trazodone , Pseudoephedrine Other mechanism.

Additional Information: Tricyclic antidepressants may increase or decrease the effect of the other drug when taken in this combination.

Trazodone + Buspirone

Trazodone and Buspirone both increase affecting serotonin levels in the blood. Too much serotonin is a potentially life-threatening situation. Severe signs and symptoms include high blood pressure and increased heart rate that lead to shock.

Trazodone + Venlafaxine

Trazodone and Venlafaxine both increase affecting serotonin levels in the blood. Too much serotonin is a potentially life-threatening situation. Severe signs and symptoms include high blood pressure and increased heart rate that lead to shock.

Trazodone + Ziprasidone

Trazodone and Ziprasidone both increase causing a dangerous abnormal heart rhythm.

Aripiprazole + Iloperidone

Aripiprazole and Iloperidone both increase causing a dangerous abnormal heart rhythm.

Aripiprazole + Ziprasidone

Aripiprazole and Ziprasidone both increase causing a dangerous abnormal heart rhythm.

Aripiprazole + Paliperidone

Aripiprazole and Paliperidone both increase causing a dangerous abnormal heart rhythm.

Aripiprazole + Trazodone

Aripiprazole and Trazodone both increase causing a dangerous abnormal heart rhythm.

Aripiprazole + Citalopram

Aripiprazole and Citalopram both increase causing a dangerous abnormal heart rhythm.

Aripiprazole + Quetiapine

Aripiprazole and Quetiapine both increase causing a dangerous abnormal heart rhythm.

Tramadol + Diphenoxylate Hcl

Tramadol , Diphenoxylate Hcl Other mechanism.

Additional Information: Tramadol may reinitiate opiate dependence in patients previously addicted to other opiates; it may also provoke withdrawal in patients who are currently opiate dependent.

Ibuprofen + Naproxen

Ibuprofen will increase the level or effect of Naproxen by acidic drugs competing for the same pathway through the kidney.

Ibuprofen and Naproxen both increase the risk of side effects and cause bleeding or bruising.

Ibuprofen and Naproxen both increase potassium levels in the blood.

Epinephrine + Ziprasidone

Epinephrine and Ziprasidone both increase causing a dangerous abnormal heart rhythm.

Escitalopram + Buspirone

Escitalopram and Buspirone both increase affecting serotonin levels in the blood. Too much serotonin is a potentially life-threatening situation. Severe signs and symptoms include high blood pressure and increased heart rate that lead to shock.

Escitalopram + Amitriptyline

Escitalopram and Amitriptyline both increase affecting serotonin levels in the blood. Too much serotonin is a potentially life-threatening situation. Severe signs and symptoms include high blood pressure and increased heart rate that lead to shock.

Escitalopram increases toxicity of Amitriptyline by causing a dangerous abnormal heart rhythm.

Escitalopram + Trazodone

Escitalopram and Trazodone both increase affecting serotonin levels in the blood. Too much serotonin is a potentially life-threatening situation. Severe signs and symptoms include high blood pressure and increased heart rate that lead to shock.

Escitalopram + Venlafaxine

Escitalopram and Venlafaxine both increase affecting serotonin levels in the blood. Too much serotonin is a potentially life-threatening situation. Severe signs and symptoms include high blood pressure and increased heart rate that lead to shock.

Escitalopram + Iloperidone

Escitalopram increases toxicity of Iloperidone by causing a dangerous abnormal heart rhythm.

Escitalopram + Paliperidone

Escitalopram increases toxicity of Paliperidone by causing a dangerous abnormal heart rhythm.

Escitalopram + Quetiapine

Escitalopram increases toxicity of Quetiapine by causing a dangerous abnormal heart rhythm.

Escitalopram + Bupropion

Escitalopram increases toxicity of Bupropion by unspecified interaction mechanism.

Additional Information: Keep bupropion dose as low as possible as seizures may occur.

Ibuprofen IV + Naproxen

Ibuprofen IV and Naproxen both increase potassium levels in the blood.

Ibuprofen IV will increase the level or effect of Naproxen by acidic drugs competing for the same pathway through the kidney.

Ibuprofen IV and Naproxen both increase the risk of side effects and cause bleeding or bruising.

MINOR POTENTIAL FOR SERIOUS NEGATIVE INTERACTIONS

These combinations of drugs have risk for dangerous interaction.

Atropine + Trazodone

Atropine and Trazodone both decrease additive side effects that may cause blurred vision, increased saliva, and increased bowel movements and urination.

Atropine increases levels of Trazodone by unknown mechanism.

Lansoprazole + Levothyroxine

Lansoprazole decreases levels of Levothyroxine by reducing stomach acidity. This interaction may occur when both drugs are taken by mouth.

Additional Information: There may or may not be an interaction between these drugs.

Lansoprazole + Lisdexamfetamine

Absorption of drugs is slowed.

Omeprazole + Lisdexamfetamine

Absorption of drugs is slowed.

Omeprazole + Clonazepam

Omeprazole increases levels of Clonazepam by slowing drug metabolism.

Omeprazole + Levothyroxine

Omeprazole decreases levels of Levothyroxine by reducing stomach acidity. This interaction may occur when both drugs are taken by mouth.

Additional Information: There may or may not be an interaction between these drugs.

Omeprazole + Sulfamethoxazole

Omeprazole will increase the level or effect of Sulfamethoxazole by altering drug metabolism.

Omeprazole + Ibuprofen IV

Omeprazole will increase the level or effect of Ibuprofen IV by altering drug metabolism.

Clonidine + Metformin

Clonidine decreases effects of Metformin by opposing drug effects.

Additional Information: Caution; use of these two drugs may lead to variations in blood glucose levels.

Guanfacine + Metformin

Additional Information: Caution; use of these two drugs may lead to variations in blood glucose levels.

Guanfacine decreases effects of Metformin by opposing drug effects.

Hydrochlorothiazide + Metformin

Hydrochlorothiazide decreases effects of Metformin by opposing drug effects.

Additional Information: Thiazide diuretics, in doses over 50mg/day, and similar drugs increase glucose levels, reducing the effect of drugs used for treating diabetes.

Hydrochlorothiazide will increase the level or effect of Metformin by nonacidic drugs competing for the same pathway through the kidney.

Hydrochlorothiazide + Sulfamethoxazole

Hydrochlorothiazide increases levels of Sulfamethoxazole by unspecified interaction mechanism.

Hydrochlorothiazide , Sulfamethoxazole Mechanism: unspecified interaction mechanism.

Additional Information: There may be an increased chance of low blood sodium (hyponatremia).

Hydrochlorothiazide will increase the level or effect of Sulfamethoxazole by nonacidic drugs competing for the same pathway through the kidney.

Hydrochlorothiazide + Ibuprofen

Hydrochlorothiazide will increase the level or effect of Ibuprofen by acidic drugs competing for the same pathway through the kidney.

Hydrochlorothiazide + Meloxicam

Hydrochlorothiazide will increase the level or effect of Meloxicam by acidic drugs competing for the same pathway through the kidney.

Hydrochlorothiazide + Naproxen

Hydrochlorothiazide will increase the level or effect of Naproxen by acidic drugs competing for the same pathway through the kidney.

Budesonide + Metformin

Budesonide decreases effects of Metformin by opposing drug effects.

Budesonide + Hydrochlorothiazide

Budesonide , Hydrochlorothiazide Mechanism: additive drug effects.

Additional Information: There may be an increased chance of low blood potassium.

Budesonide + Ziprasidone

Budesonide will decrease the level or effect of Ziprasidone by altering drug metabolism.

Budesonide + Amitriptyline

Budesonide will decrease the level or effect of Amitriptyline by altering drug metabolism.

Budesonide + Oxybutynin

Budesonide will decrease the level or effect of Oxybutynin by altering drug metabolism.

Budesonide + Montelukast

Budesonide will decrease the level or effect of Montelukast by altering drug metabolism.

Clonazepam + Acetaminophen

Clonazepam decreases levels of Acetaminophen by speeding up drug metabolism.

Additional Information: May cause liver problems.

Lamotrigine + Acetaminophen

Lamotrigine decreases levels of Acetaminophen by speeding up drug metabolism.

Additional Information: May cause liver problems.

Levetiracetam + Acetaminophen

Levetiracetam decreases levels of Acetaminophen by speeding up drug metabolism.

Additional Information: May cause liver problems.

Topiramate + Acetaminophen

Topiramate decreases levels of Acetaminophen by speeding up drug metabolism.

Additional Information: May cause liver problems.

Topiramate + Montelukast

Topiramate will decrease the level or effect of Montelukast by altering drug metabolism.

Topiramate + Oxybutynin

Topiramate will decrease the level or effect of Oxybutynin by altering drug metabolism.

Topiramate + Ziprasidone

Topiramate will decrease the level or effect of Ziprasidone by altering drug metabolism.

Oxybutynin + Acetaminophen

Oxybutynin decreases levels of Acetaminophen by unspecified interaction mechanism.

Fluoxetine + Lansoprazole

Fluoxetine will increase the level or effect of Lansoprazole by altering drug metabolism.

Fluoxetine + Omeprazole

Fluoxetine will increase the level or effect of Omeprazole by altering drug metabolism.

Venlafaxine + Fluoxetine

Venlafaxine will increase the level or effect of Fluoxetine by altering drug metabolism.

Amitriptyline + Metformin

Amitriptyline increases effects of Metformin by added drug effects.

Amitriptyline + Atropine

Amitriptyline increases levels of Atropine by unknown mechanism.

Trazodone + Metformin

Trazodone increases effects of Metformin by added drug effects.

Trazodone + Fexofenadine

Trazodone will decrease the level or effect of Fexofenadine by affects how the drug is eliminated from the body (via what is known as the P-glycoprotein [MDR1] transporter).

Trazodone + Loratadine

Trazodone will decrease the level or effect of Loratadine by affects how the drug is eliminated from the body (via what is known as the P-glycoprotein [MDR1] transporter).

Cephalexin + Meloxicam

Cephalexin will increase the level or effect of Meloxicam by acidic drugs competing for the same pathway through the kidney.

Cephalexin + Hydrochlorothiazide

Cephalexin will increase the level or effect of Hydrochlorothiazide by acidic drugs competing for the same pathway through the kidney.

Cephalexin + Naproxen

Cephalexin will increase the level or effect of Naproxen by acidic drugs competing for the same pathway through the kidney.

Cephalexin + Ibuprofen

Cephalexin will increase the level or effect of Ibuprofen by acidic drugs competing for the same pathway through the kidney.

Ciprofloxacin + Clonazepam

Ciprofloxacin increases levels of Clonazepam by slowing drug metabolism.

Sulfamethoxazole + Amitriptyline

Sulfamethoxazole decreases levels of Amitriptyline by unspecified interaction mechanism.

Sulfamethoxazole + Trazodone

Sulfamethoxazole decreases levels of Trazodone by unspecified interaction mechanism.

Sulfamethoxazole + Meloxicam

Sulfamethoxazole will increase the level or effect of Meloxicam by altering drug metabolism.

Sulfamethoxazole + Ibuprofen IV

Sulfamethoxazole will increase the level or effect of Ibuprofen IV by altering drug metabolism.

Sulfamethoxazole + Ibuprofen

Sulfamethoxazole will increase the level or effect of Ibuprofen by altering drug metabolism.

Sulfamethoxazole + Metformin

Sulfamethoxazole will increase the level or effect of Metformin by nonacidic drugs competing for the same pathway through the kidney.

Ibuprofen + Meloxicam

Ibuprofen will increase the level or effect of Meloxicam by acidic drugs competing for the same pathway through the kidney.

Meloxicam + Naproxen

Meloxicam will increase the level or effect of Naproxen by acidic drugs competing for the same pathway through the kidney.

Albuterol + Hydrochlorothiazide

Albuterol , Hydrochlorothiazide Mechanism: additive drug effects.

Additional Information: Low potassium (hypokalemia).

Epinephrine + Hydrochlorothiazide

Epinephrine , Hydrochlorothiazide Mechanism: additive drug effects.

Additional Information: Low potassium (hypokalemia).

Dexmethylphenidate + Amitriptyline

Dexmethylphenidate increases effects of Amitriptyline by slowing drug metabolism.

Dexmethylphenidate + Trazodone

Dexmethylphenidate increases effects of Trazodone by slowing drug metabolism.

Dexmethylphenidate + Clonazepam

Dexmethylphenidate increases effects of Clonazepam by slowing drug metabolism.

Dexmethylphenidate + Gabapentin

Dexmethylphenidate increases effects of Gabapentin by slowing drug metabolism.

Dexmethylphenidate + Lamotrigine

Dexmethylphenidate increases effects of Lamotrigine by slowing drug metabolism.

Dexmethylphenidate + Levetiracetam

Dexmethylphenidate increases effects of Levetiracetam by slowing drug metabolism.

Dexmethylphenidate + Topiramate

Dexmethylphenidate increases effects of Topiramate by slowing drug metabolism.

Dexmethylphenidate + Fluoxetine

Dexmethylphenidate increases effects of Fluoxetine by slowing drug metabolism.

Dexmethylphenidate + Venlafaxine

Dexmethylphenidate increases effects of Venlafaxine by slowing drug metabolism.

Dextroamphetamine + Diphenoxylate Hcl

Dextroamphetamine increases effects of Diphenoxylate Hcl by unspecified interaction mechanism.

Dextroamphetamine + Tramadol

Dextroamphetamine increases effects of Tramadol by unspecified interaction mechanism.

Loratadine + Fexofenadine

Loratadine will increase the level or effect of Fexofenadine by affects how the drug is eliminated from the body (via what is known as the P-glycoprotein [MDR1] transporter).

Amantadine + Oxybutynin

Amantadine increases effects of Oxybutynin by added drug effects.

Amantadine + Lisdexamfetamine

Amantadine , Lisdexamfetamine . Either increases the level of the other by added drug effects.

Additional Information: Combination increases stimulation of the nervous system.

Amantadine + Dexmethylphenidate

Amantadine , Dexmethylphenidate . Either increases the level of the other by added drug effects.

Additional Information: Combination increases stimulation of the nervous system.

Amantadine + Dextroamphetamine

Amantadine , Dextroamphetamine . Either increases the level of the other by added drug effects.

Additional Information: Combination increases stimulation of the nervous system.

Amantadine + Methylphenidate

Amantadine , Methylphenidate . Either increases the level of the other by added drug effects.

Additional Information: Combination increases stimulation of the nervous system.

Escitalopram + Labetalol

Escitalopram increases levels of Labetalol by slowing drug metabolism.

Ethinylestradiol + Amitriptyline

Ethinylestradiol , Amitriptyline Mechanism: unspecified interaction mechanism.

Additional Information: Estrogens and progestins may decrease effectiveness of tricyclic antidepressant drugs.

Ethinylestradiol + Trazodone

Ethinylestradiol , Trazodone Mechanism: unspecified interaction mechanism.

Additional Information: Potential for increase in antidepressant side effects.

Benztropine + Trazodone

Benztropine and Trazodone both decrease additive side effects that may cause blurred vision, increased saliva, and increased bowel movements and urination.

Ibuprofen IV + Meloxicam

Ibuprofen IV will increase the level or effect of Meloxicam by acidic drugs competing for the same pathway through the kidney.